

# Family-based cognitive factors effective on preventing the onset of substance use in Iranian Society's Children: applying the intervention mapping protocol

Citation for published version (APA):

Mirzaei-Alavijeh, M., Alirezahidarnia, Kok, G., Niknami, S., & Motlagh, M. E. (2016). Family-based cognitive factors effective on preventing the onset of substance use in Iranian Society's Children: applying the intervention mapping protocol. *Acta Medica Mediterranea*, 32, 1015-1020.  
<http://www.actamedicamediterranea.com/archive/2016/special-issue-2/family-based-cognitive-factors-effective-on-preventing-the-onset-of-substanceuse-in-iranian-societys-children-applying-the-intervention-mapping-protocol>

## Document status and date:

Published: 01/01/2016

## Document Version:

Publisher's PDF, also known as Version of record

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

Download date: 05 May. 2023

## FAMILY-BASED COGNITIVE FACTORS EFFECTIVE ON PREVENTING THE ONSET OF SUBSTANCE USE IN IRANIAN SOCIETY'S CHILDREN: APPLYING THE INTERVENTION MAPPING PROTOCOL

MEHDI MIRZAEI-ALAVIJEH<sup>1</sup>, ALIREZAHIDARNIA<sup>1\*</sup>, GERJO KOK<sup>2</sup>, SHAMSADDIN NIKNAMI<sup>1</sup>, MOHAMMAD ESMAIL MOTLAGH<sup>3</sup>

<sup>1</sup>Department of Health Education, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran - <sup>2</sup>Department of Work and Social Psychology School of Psychology and Neuroscience, Maastricht University, Maastricht, Netherlands - <sup>3</sup>Department of Pediatrics, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

### ABSTRACT

**Introduction:** According to increase prevalence rate of substance abuse and appears a reduction in mean age of onset substance use in Iranian society and importance of developing theory and evidence-based prevention family skills training programs the present study was done aimed to investigate family-based cognitive factors effective in preventing the onset of substance use in Iranian society's children by applying the intervention mapping protocol.

**Materials and methods:** This cross-sectional study was carried out among 234 Iranian mothers in Tehran who were randomly selected to participate voluntarily in the study. The participants filled out a self-administered questionnaire. Data were analyzed by the SPSS software (ver. 21.0) using Pearson correlations, independent t-test, ANOVA and linear regression at 95% significant level.

**Results:** Knowledge, attitude, perceived risk, subjective norms, and self-efficacy were cognitive factors that accounted for 48% of the variation in the outcome measure of the behaviors in preventing the onset of substance use among children. Linear regression showed that self-efficacy ( $\beta=0.509$  &  $P=0.001$ ), attitude ( $\beta=0.436$  &  $P=0.009$ ), subjective norms ( $\beta=0.344$  &  $P=0.028$ ) and Behavioural intention ( $\beta=0.222$  &  $P=0.001$ ) were the most influential predictors on preventing the onset of substance use among children.

**Conclusion:** The findings suggest that to developing a theory and evidence-based prevention family skills training program it will be useful to increase mothers' self-efficacy, attitude, subjective norms, and Behavioural intention to preventing the onset of substance use among children.

**Key words:** Family Skills Training Program, Substance Abuse, Cognitive Factors, Parents.

Received February 05, 2016; Accepted March 02, 2016

### Introduction

During the recent years, drug abuse has been widely reported in Iran. Studies since 2000 to 2009 suggested that, after inflation and unemployment, drug abuse is the third major problem in Iran<sup>(1)</sup>. Also, drug abuse age has significantly decreased<sup>(2)</sup>. Thus, it seems necessary to prevent substance use since childhood. However genetics, natural and environmental factors all affect a child development, family dynamics plays an essential role in child development<sup>(3)</sup>.

Also, researchers investigated various models to explain different factors influence the possibility to substance abuse and reported parental and family

factors have a central position in the long-term pathways leading to substance abuse, while peers effect was a often the major reason adolescents initiate drug use in lower ages. In other words, a positive family environment is the primary reason youth do not engage in risky behaviors such as substance abuse<sup>(4)</sup>. Studies listed the following effective family factors to protect children from the onset of substance use: secure and healthy parent/child attachment; parental supervision, monitoring and effective discipline; Communication of pro-social family values; Parental involvement in child's life; and Supportive parenting (emotionally, cognitively, socially and financially)<sup>(5)</sup>.

Furthermore, studies suggested evidence on certain family features which have been known as major risk factors, such as poor management on children behavior, rough and contradictory disciplines, lack of opportunity to learn social skills to resist against social and educational problems teenagers may face in society<sup>(6)</sup>.

Generally, a chaotic family environment and lack of organization in family have been known as risk factors to substance abuse<sup>(7)</sup>. In addition, apathetic, non-responsive parent-child relationship, emotional insecurity and lack of organized care and control from parents to provide peace to children during their first years of growth increase the risk of anxiety, depression and communication problems among children and young adults<sup>(8)</sup>.

Studies on the field showed that major family factors which increase the possibility from the onset of substance use among children include: Lack of bonding and insecure relationship with parents; lack of a significant relationship with a caring adult; ineffective parenting; chaotic home environment; Parents or siblings who abuse substances, suffer from mental illness or are involved in criminal behavior; antisocial isolation<sup>(8)</sup>.

Recent research reported family skill training including: parents active participation, developing social skills and accountability among children, and considering substance abuse issues, to be the most influential program to prevent the onset of substance use at low ages<sup>(9)</sup>.

In present study to identify and prioritize risky and protective family based factors to the onset of substance use among children and also parents cognitive determinants in this regards, we used the first and the second steps of the IM protocol in the presence of a planning group, including stakeholders related to family-based substance abuse preventions among children<sup>(9)</sup>.

Based on intervention mapping protocol, in first step by using of relevant literature, evidence and collective planning group wisdom identified effective family based preventing factors in the onset of substance use among children and also addressed to prioritized the most important and the most changeable factors (according to the literature and evidence) at a sample of families selective according to cultural factors in the society under study<sup>(10)</sup>. Since the awareness of current conditions related to cognitive determinants about effective family-based risky and protective factors the onset of substance use among children in under study society is essen-

tial to developing preventive programs, the present study was done aims to investigate family based cognitive factors to prevent the onset of substance use among children based on intervention mapping protocol.

## Material and methods

It was a cross-sectional study that carried out among 234 randomly selected mothers in Tehran, Iran. Considering the results from the pilot study phase and dependent variables  $SD=5.35$  related to mothers behavior about preventing the onset of substance use among children, at 95% significant level and  $d=0.5$ , the number of samples was defined to be 229 and considering 10% attrition rate, a sample of 252 was estimated. A questionnaire was developed to the goal of the study, based on evidence of the previous theory based IM step two and studies on substance abuse prevention to provide determinants items include: knowledge, attitude, perceived risk, subjective norms, self-efficacy, behavioral intentions and actual behaviors to collect required data. Participants were asked to choose their answers from a 5-Likert scale ranging from completely agree to completely disagree.

Background data gathered included age (year), educational level (under diploma (12 grades <), Diploma (12 grades), and academic), family and friends history on substance abuse (yes, no), occupation (housewife, working), and family economic status (very good, good, average, poor, very poor). seven determinants include: four items on attitude ( $\alpha=0.70$ ) scores ranging from 5 to 20 (e.g. "I believe that family awareness of each other's feelings, interests, beliefs, opinions, experiences, goals and decision is effective on preventing the onset of substance use among my children"), six items on perceived risk ( $\alpha=0.83$ ) scores ranging from 6 to 30 (e.g. "Defining and discussing family values and expectations about substance use have no effect on the onset of substance use among my children"), one item on knowledge, scores ranging from 1 to 5 (Family contention and not discussing to find the solutions is effective on the onset of substance use among children), eight items on self-efficacy ( $\alpha=0.79$ ) scores ranging from 8 to 40 (e.g. "I have the skills to avoid disagreement to my spouse and find proper solutions through discussions to overcome the difficulties made to prevent of the onset of substance use among my children"), two items on subjective norms ( $\alpha=0.86$ ) scores ranging from 2 to 10 (e.g.

“Praising positive behaviors (for instance; paying attention to family and social values and rules, and taking responsibilities) is not important in my family to prevent the onset of substance use among my children”), behavioral intension with ten items ( $\alpha=0.74$ ) scores ranging from 10 to 50 (e.g. “I tend clarify substance use regulations in my family to prevent the onset of substance use among my children”), and behavior with 10 items ( $\alpha=0.81$ ) scores ranging from 10 to 50 (e.g. “I spend more time with my family and I responsible to my family members requirements to prevent the onset of substance use among my children”). To confirm content validity index (CVI) and content validity ratio (CVR) of the questionnaire were used through consulting experts in health educating, addiction, family and psychology and face validity was approved based on the qualitatively collected view points of a target group, similar to the under study population, about difficulty, appropriateness and ambiguity of items. Inclusion and exclusion criteria to the study were: healthy family framework, being Tehran resident, age range 30 to 50, having a child at pre-school level, no substance use in parents or other family members. Self-reports were the bases to confirm non substance user. Also all the participants were justified on how to study and confidentiality of information as well as the aim of this study and directed to the study on their own willingness in order to comply with research ethics. The collected data were analyzed using SPSS version 21 making use of descriptive and analytical statistics: Pearson correlation, independent t-test, ANOVA and linear regression analysis at 95% significant level.

## Results

The mean age of respondents was 35.92 years [SD: 4.03], ranged from 30 to 45 years. Demographic characteristics of the participants are shown in Table 1. There was a significant correlation between having family and friends history of substance abuse, educational level and behaviors' preventing in onset of substance use in Iranian society's children. However, there was no significant difference between behaviors' preventing in onset of substance use and job, economic status (Table 1). In addition, Bivariate associations among the age, and behaviors' preventing in onset of substance use showed statistically not significant ( $r=0.124$  &  $P=0.059$ ).

Variable		n (%)	Mean	SD	P-value
Education level	Under Diploma (12 grades<)	19 (8.1 %)	31.31	4.53	0.028
	Diploma (12 grades)	179 (76.5 %)	31.88	4.41	
	Academic	36 (15.4 %)	34.08	6.04	
Economic Status	Very Poor	13 (5.6 %)	29.84	3.43	0.239
	Poor	23 (9.8 %)	31.86	4.72	
	Average	143 (61.1 %)	32.63	4.63	
	Good	45 (19.2 %)	31.48	5.08	
	Very Good	10 (4.3 %)	32.4	6.05	
Positive Family history of Substance	Yes	4 (1.7 %)	31.25	7.88	0.696
Abuse	No	230 (98.3 %)	32.19	4.71	
Positive Friends history of Substance	Yes	5 (2.1 %)	30.8	6.22	0.515
Abuse	No	229 (97.9 %)	32.2	4.73	
Occupation	Housewife	212 (90.6 %)	32.07	4.63	0.298
	Working	22 (9.4 %)	33.18	5.85	

**Table 1:** Association between background variable and behaviors' preventing the onset of substance use among children.

	Mean (SD)	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	X <sup>5</sup>	X <sup>6</sup>
X <sup>1</sup> . Attitude	15.76 (1.75)	1					
X <sup>2</sup> . Perceived Risk	22.36 (2.44)	0.321**	1				
X <sup>3</sup> . Knowledge	3.68 (0.52)	0.251**	0.280**	1			
X <sup>4</sup> . Self-Efficacy	28.26 (3.49)	0.600**	0.285**	0.208**	1		
X <sup>5</sup> . Subjective Norms	6.66 (1.62)	0.306**	0.433**	0.343**	0.235**	1	
X <sup>6</sup> . Behavioural Intention	32.72 (5.32)	0.397**	0.352**	0.380**	0.447**	0.423**	1
X <sup>6</sup> . Behaviour	32.17 (4.75)	0.520**	0.330**	0.286**	0.609**	0.360**	0.529**

**Table 2:** Predictor variables of cognitive factors effective on preventing the onset of substance use among children.

Variable	B	SE B	Beta	T	P-value
<b>Step 1</b>					
Attitude	0.413	0.167	0.152	2.468	0.014
Perceived Risk	0.079	0.107	0.041	0.737	0.462
Knowledge	0.336	0.479	0.037	0.701	0.484
Self-Efficacy	0.505	0.085	0.371	5.968	0.001
Subjective Norms	0.284	0.166	0.097	1.707	0.089
Behavioural intention	0.209	0.053	0.234	3.939	0.001
<b>Step 2</b>					
Attitude	0.421	0.167	0.155	2.524	0.012
Perceived Risk	0.086	0.106	0.044	0.805	0.421
Self-Efficacy	0.504	0.085	0.370	5.964	0.001
Subjective Norms	0.303	0.164	0.103	1.849	0.066
Behavioural intention	0.217	0.052	0.243	4.205	0.001
<b>Step 3</b>					
Attitude	0.436	0.166	0.161	2.632	0.009
Self-Efficacy	0.509	0.084	0.373	6.034	0.001
Subjective Norms	0.344	0.155	0.118	2.215	0.028
Behavioural intention	0.222	0.051	0.249	4.345	0.001
SE=Standard Error					
Final model: Step 3, Adjusted R squared=0.48, F= 53.938, and P<0.001					

**Table 3:** Predictors of mother's behaviors preventing the onset of substance use among children based on linear regression analysis.

Table 2 shows bivariate correlations between the predictor variables of cognitive factors effective in based on bivariate correlation analysis. Finally, linear regression analysis was performed to explain the variation of mother's behaviors preventing the onset of substance use among children. As shown in Table 3, collectively, Predictors variables accounted for 48% of the variation seen in behaviors preventing the onset of substance use in Iranian society's children.

## Discussion

Our findings indicated that the attitude, subjective norms, self-efficacy, and behavior intention were the three main cognitive factors which were effective in preventing the onset of substance use in Iranian society's children. Many studies have addressed to prediction of substance use behavior by using cognitive construct, models, and theories<sup>(12)</sup>. In this regard, Mirzaei-Alavijeh et al. in their study on role of fathers in prevention of children tendency toward addictive drugs based on theory of planned behavior indicated subjective norms and behavior intention were the best predictors for fathers' behavior about prevention of children tendency toward addictive drugs<sup>(13)</sup>.

In addition van De Ven<sup>(14)</sup>, Harakehet al<sup>(15)</sup> reported behavioral intention were as important predictors for onset of cigarette smoking among adolescents. Jalilian et al. presented attitude, outcome expectancies, and subjective norms were the most influential predictors for drug abuse among a sample of Iranian male medical college students<sup>(16)</sup>. Mirzaei-Alavijeh et al. carried out a research based on prototype willingness model among 18 to 35 years old college students in the center of Iran their results showed that attitude, subjective norms and risk images were the best predictors in preventing of drug abuse<sup>(17)</sup>. In addition, Gerrard et al. reported the high correlation between behavior intention and drug abuse<sup>(18)</sup>. Also, Eslamiet al. showed that behavioral intention was a stronger prediction factor to Ritalin misuse among Iranian medical college students.

Since the present study investigated behavior intention construct as intention performance rather than time of intention, this investigation under the supervision of the experts in the field could be a better predictor to predict behavior onset speed and possible actual behavior<sup>(20)</sup>. Therefore, considering the role of behavioral intention to predict behavior

of preventing the onset of substance use among children in the present study, it seems this behavioral intention construct could be effective on behavior change in intervention program. Based on the our results from the present study and other related studies, cognitive factors, particularly attitude and self-efficacy, play important roles to predict substance abuse and its prevention behaviors, therefore it is essential to consider cognitive determinants to provide preventive intervention programs.

Another finding of this study indicated cognitive factors mentioned accounted for 48% of the variation of parent's behaviors on preventing the onset of substance use among Iranian society's children. Many studies have addressed the cognitive factors their results accounted between 17 to 63% of the variation in the outcome measure of the behavior in substance abuse and substance use preventing. For example Mirzaei-Alavijeh et al. showed that behavioral intention predicted 17.9% of the variance of father behaviors on prevention of children tendency toward addictive drugs in Islamic republic of Iran<sup>(21)</sup>.

Eslamiet al. showed that three predictor variables include; attitude, subjective norms, and prototype accounted for 29% of the variation in intention to Ritalin misuse among Iranian medical college students<sup>(19)</sup>. Mirzaei-Alavijeh et al. showed that theory of planned behavior accounted for 23.5 % of the variance of father behaviors on prevention of children tendency toward addictive drugs in Islamic republic of Iran<sup>(13)</sup>. Also, Van De Ven et al<sup>(14)</sup> and Harakehet al<sup>(15)</sup> reported for 16% and 38%, respectively of the variation in the outcome measure of the intention to the onset of among parents and adolescents. As well as, Jalilian et al. presented attitude, outcome expectation, outcome expectancies, subjective norms, and self-control were cognitive factors that accounted for 49% of the variation in the outcome measure of the intention to drugs abuse among a sample of Iranian male medical college students<sup>(16)</sup>.

This difference in prediction amount could be relative to evaluation of outcome construct; individuals; and social, cultural bases under study. Our results showed that among underlying variables under study, there was a significant relationship between educational level and preventive behaviors, so that the higher educated the mothers, had the more appropriate behaviors to prevent the onset of substance use in their children.

Results from A potential limitation is that the behavioral and psychological questionnaires used in

the study rely on thyself-report format. The results, therefore, may be subject to self-report bias, resulting in some unspecified amount of misclassification as to the participants' actual levels of preventive behavior about the onset of substance use among children. However, population-based studies must rely on self-report to a large extent because there is frequently no direct contact between the subject and the investigator. The accuracy of self-report has been extensively investigated in the area of health-related behaviors, where it has been found to be accurate<sup>(22)</sup>. Also another limitation was lack of relative data on fathers.

## References

- 1) Moghanibashi-Mansourieh, Amir, and Abbas Deilamizade. *The state of data collection on addiction in Iran*. Addiction 2014; 109(5): 854-854.
- 2) United Nations Office on Drugs and Crime (UNODC). *World Drug Report*. New York: UNODC; 2013.
- 3) Glantz MD, Leshner AI. Drug abuse and developmental psychopathology. *Development and Psychopathology* 2000; 12(04): 795-814.
- 4) Kumpfer KL, Alvarado R, Whiteside HO. *Family-based interventions for substance use and misuse prevention*. Substance use & misuse 2003; 38(11-13): 1759-1787.
- 5) Sale E, Sambrano S, Springer JF, Turner, CW. *Risk, protection, and substance use in adolescents: a multi-site model*. Journal of drug education 2003; 33(1), 91.
- 6) Forgatch MS, Knutson NM, Nancy M. *Linking basic and applied research in a prevention science process*. Liddle HA; Santisteban DA; Levant RF, Bray JH. *Family psychology: Science-based interventions*. Washington, DC, US: American 2002: 239-257.
- 7) United States of America, Department of Health and Human Services, National Institutes of Health, Preventing Drug Use Among Children and Adolescents: a Research-Based Guide for Parents, Educators, and Community Leaders, 2nd ed, NIH publication No. 04-4212(A), Bethesda, Maryland, National Institute on Drug Abuse 2003.
- 8) Porter LS, Porter BO. *A blended infant massage-parenting enhancement program for recovering substance-abusing mothers*. Pediatric Nursing 2004; 30(5): 363-401.
- 9) Petrie J, Bunn F, Byrne G. *Parenting programmes for preventing tobacco, alcohol or drugs misuse in children < 18: a systematic review*. Health education research 2007; 22(2): 177-191.
- 10) Kok G. *A practical guide to effective behavior Change: How to apply theory- and evidence-based behavior change methods in an intervention*. The European Health Psychologist 2014; 16(5): 156- 170.
- 11) Mirzaei-Alavijeh M, Jalilian F, Zinat Motlagh F, Mazloomi Mahmodabad S, Zolghadr R, Hatamzadeh N. *Effectiveness of Drug Abuse Preventative Intervention among Iranian Medical College Students Based on the Theory of Planned Behavior*. Health Education & Health Promotion 2015; 2(1): 41-52.
- 12) Jalilian F, Karami-Matin B, Ahmadpanah M, Ataee M, AhmadiJouybari T, Eslami AA, Mirzaei-Alavijeh M. *Socio-demographic characteristics associated with cigarettes smoking, drug abuse and alcohol drinking among male medical university students in Iran*. Journal of research in health sciences. 2015 25; 15 (1): 42-46.
- 13) Mirzaei-Alavijeh M, Mazloomi S, Yassini M, Askarshahi M, Jalilian F. *The Role of Fathers in prevention of Children Tendency toward Addictive Drugs: an Application of Theory of Planned Behavior*. Journal of Police Medicine 2015; 3(4): 193-202. (Persian).
- 14) Van De Ven MO, Engels RC, Otten R, Van Den Eijnden RJ. *A longitudinal test of the theory of planned behavior predicting smoking onset among asthmatic and non-asthmatic adolescents*. Journal of behavioral medicine 2007; 30(5): 435-45.
- 15) Harakeh Z, Scholte RH, Vermulst AA, de VH, Engels RC. *Parental factors and adolescents' smoking behavior: an extension of the theory of planned behavior*. Preventive medicine 2004; 39(5): 951-61.
- 16) Jalilian F, Ataee M, Matin BK, Ahmadpanah M, Jouybari TA, Eslami AA, Mahboubi M, Mirzaei-Alavijeh M. *Cognitive factors related to drug abuse among a sample of Iranian male medical college students*. Global journal of health science. 2015; 7(5): 143-51.
- 17) Mirzaei-Alavijeh M, Jalilian F, Movahed E, Mazloomi S, ZinatMotlagh F, Hatamzadeh, N. *Predictors of drug abuse among students with application of Prototype/Willingness Model*. Journal of Police Medicine 2013; 2(2), 111-118.
- 18) Gerrard M, Gibbons FX, Brody GH, Murry VM, Cleveland MJ, Wills TA. *A theory-based dual-focus alcohol intervention for preadolescents: The Strong African American Families Program*. Psychology of Addictive Behaviors 2006; 20, 185-95.
- 19) Eslami AA, JalilianF,Ataee M, Mirzaei-Alavijeh M, Mahboubi M, Afsar A, Aghaei A. *Intention and willingness in understanding Ritalin misuse among Iranian medical college students: a cross-sectional study*. Global journal of health science 2014; 6(6): 43-53.
- 20) Ajzen I. *The theory of planned behavior*. Organizational Behavior and Human Decision Processes, 1991; 50(2): 179-211.
- 21) Mirzaei-Alavijeh M, Mazloomi S, Yassini M, Askarshahi M, JalilianF, Zinat Motlagh F, Hatamzadeh N. *Fathers' behavioral intention and behavior in prevention of children tendency toward addictive drugs*. Iranian Journal of Health Education and Health Promotion 2013; 1(2): 57-66. (Persian).
- 22) Velicer WF, Prochaska JO, Rossi JS, Snow MG. *Assessing outcome in smoking cessation studies*. Psychological bulletin 1992; 111(1): 23-41.

*Acknowledgements*

*This article is a part of the Ph.D. dissertation of health education and health promotion supported by TarbiatModares University. We would like to thank Deputy of Research of TarbiatModares University for financial support.*

---

*Corresponding author*  
hidarnia@modares.ac.ir